

Comments on conformal higher spin theory

Arkady Tseytlin

Based on:

“On triviality of S-matrix in conformal higher spin theory”

with M. Beccaria and S. Nakach [arXiv:1607.06379](https://arxiv.org/abs/1607.06379)

“On conformal higher spins in curved background”

with M. Grigoriev [arXiv:1609.09381](https://arxiv.org/abs/1609.09381)

“On induced action for conformal higher spins in curved background”

with M. Beccaria [arXiv:1702.00222](https://arxiv.org/abs/1702.00222)

- free complex scalar: $\square\Phi = 0$

conserved $J_\mu = i(\Phi^*\partial_\mu\Phi - \partial_\mu\Phi^*\Phi)$ and stress $T_{\mu\nu}$

couple to external sources

$$L = \partial^\mu\Phi^*\partial_\mu\Phi + A^\mu(x)J_\mu + h^{\mu\nu}(x)T_{\mu\nu} + \dots$$

integrate out Φ : local ($\log \infty$) part of 1-loop effective action

induced Maxwell + Weyl theory

$$S = \int d^4x(-F_{\mu\nu}^2 + C_{\mu\nu\lambda\rho}^2)$$

- free scalar equation admits also higher conserved currents:

$$J_{\mu_1\dots\mu_s} = \Phi^*\partial_{\mu_1}\dots\partial_{\mu_s}\Phi + \dots, \quad s = 1, 2, 3, \dots$$

charges \rightarrow infinite dim global symmetry

corresponding sources $h_{\mu_1\dots\mu_s}$ – symmetric traceless tensors:

conformal higher spins (CHS)

- induced action for infinite tower of fields

generalizes Maxwell and Weyl: $S = \int d^4x \sum_s h_s \partial^{2s} h_s + \dots$

- local action with symmetry $\delta h_s = \partial\epsilon_{s-1} + \eta_2\alpha_{s-2} + \dots$

Motivation to study:

- unusual properties and simplifications due to underlying infinite-dimensional conformal HS symmetry (sums over infinite set of HS contributions, regularization consistent with symmetry)
- close connection to massless HS fields in AdS

CHS as toy model to study implications of HS symmetry:

- trivial partition function on a sphere
 - trivial near-flat-space S-matrix (cf. Coleman-Mandula)
 - cancellation of conformal anomalies
- fundamental role of local conformal invariance?
existence of consistent (UV finite, anomaly free)
conformal higher spin theories? unitary issue?

Plan:

- flat space background:

action for CHS as induced one and S-matrix

- curved space background:

curved space CHS operators, partition function on S^4 ,

a and c Weyl anomaly coefficients

Consistent HS theories:

- massless HS theory in AdS_{d+1} :

2-derivative kin term (unitary) but non-flat vacuum
dual to free CFT_d : e.g. scalar in vector rep of $U(N)$

S-matrix is “simple”:

reproduces correlators of currents in free CFT

- conformal HS theory:

higher derivative kin term (non-unitary) but admits flat vacuum

S-matrix is “trivial” after summation over all spin exchanges

consistent with HS symmetry

Conformal higher spin theory ($d = 4$)

- generalization of Maxwell and Weyl:

$$F_{\mu\nu}^2 \sim h_1 \partial^2 h_1, \quad C_{\mu\nu\kappa\lambda}^2 \sim h_2 \partial^4 h_2 + \partial^4 h_2 h_2 h_2 + \dots$$

- differential + algebraic (“Weyl”) gauge symmetry

$$\delta h_s = \partial \epsilon_{s-1} + \eta_2 \alpha_{s-2}$$

can gauge-fix h_s to be transverse **and** traceless off-shell

- totally symmetric $h_{\mu_1 \dots \mu_s}$ describes “pure” spin s :

maximal gauge symm consistent with locality at expense of

higher-derivative kin terms [Fradkin, AT 85]

$$S_s^{(0)} = \int d^4x h_s P_s \partial^{2s} h_s$$

$P_s \sim (\delta_{\nu}^{\mu} - \frac{\partial^{\mu} \partial_{\nu}}{\partial^2})^s$ – transv. traceless projector

- $\Delta(h_s) = 2 - s$: dimensionless coupling const
- interacting action consistent with symmetries can be defined as local induced action from scalar loop

- conformally invariant in flat space

with number of derivatives in vertices fixed by dimensions

$$S_s = \frac{1}{g^2} \sum_s \int d^4x \left(h_s \partial^{2s} h_s + \partial^{s_1+s_2+s_3-2} h_{s_1} h_{s_2} h_{s_3} \right. \\ \left. + \partial^{s_1+s_2+s_3+s_4-4} h_{s_1} h_{s_2} h_{s_3} h_{s_4} + \dots \right)$$

- conformal symmetry: CHS can be consistently defined on any conformally flat background
- admits a background-independent formulation and in general consistently defined near any curved Bach-flat (e.g. Ricci-flat) background

Properties of free CHS theory

- regularized total number of d.o.f. =0:

$$\nu_{\text{tot}} = \sum_{s=0}^{\infty} \nu_s = 0, \quad \nu_s = s(s+1) = 2, 6, \dots$$

regularization: $\sum_{s=0}^{\infty} f(s) \rightarrow \sum_{s=0}^{\infty} f(s) e^{-\epsilon(s+\frac{1}{2})} \Big|_{\text{fin.}}$

- equivalently, flat-space partition function is trivial:

$$Z_s = \left[\frac{(\det \square_{s-1})^{s+1}}{(\det \square_s)^s} \right]^{1/2} = (Z_0)^{\nu_s}, \quad Z_0 = (\det \Delta_0)^{-1/2}$$

$$Z = \prod_{s=0}^{\infty} (Z_0)^{\nu_s} = (Z_0)^{\nu_{\text{tot}}} = 1$$

- with same regularization: $Z_{\text{CHS}}(S^4)=1, \quad \sum_{s=1}^{\infty} a_s = 0$

consistent with relation between

1-loop Z of massless HS in AdS_5 and Z of CHS on S^4

[Giombi, Klebanov, Pufu, Safdi, Tarnopolsky 13; AT 13; Beccaria, Bekaert, AT 14]

- this definition of \sum_s should be consistent with underlying HS symmetry of CHS theory

“Quantized particle” approach: symmetries

start with quantized particle in external fields [Segal 02]

general phase space Hamiltonian $H(x, p)$

$$H(x, p) = \sum_{s=0}^{\infty} h^{\mu_1 \dots \mu_s}(x) p_{\mu_1} \dots p_{\mu_s} = h_0(x) + h^{\mu\nu}(x) p_{\mu} p_{\nu} + \dots$$

*-product of Weyl symbols \rightarrow product of operators

$$* = \exp \left[\frac{1}{2} \left(\overleftarrow{\partial} \frac{\partial}{\partial x^{\mu}} \frac{\partial}{\partial p_{\mu}} - \frac{\overleftarrow{\partial}}{\partial p_{\mu}} \frac{\partial}{\partial x^{\mu}} \right) \right]$$

Symmetries: canonical transfs of constraint $H(x, p) = 0$

$$\delta H = [H, \epsilon(x, p)]_* + \{H, \alpha(x, p)\}_*$$

gradient ϵ and algebraic α

- Quantum theory - in x representation: $\hat{H}\Phi(x) = 0$
action for scalar field in non-trivial background $\{h_s\}$:

$$\mathcal{S}[\phi, H] = \int d^4x \Phi^*(x) \hat{H}(x, \partial_x) \Phi(x)$$

- Invariant under the gauge transformations of Φ and h_s

$$\delta\Phi = -(\hat{\epsilon} + \hat{\alpha})\Phi, \quad \delta H = [H, \epsilon(x, p)]_* + \{H, \alpha(x, p)\}_*$$

- Choice of vacuum expansion point:

$$H = H_{\text{vac}} + h(x, p), \quad h(x, p) = \sum_{s=0}^{\infty} h^{\mu_1 \dots \mu_s}(x) p_{\mu_1} \dots p_{\mu_s}$$

$$\mathcal{S} = \int d^4x \left[\Phi^*(x) \hat{H}_{\text{vac}} \Phi(x) + \sum_s h^{\mu_1 \dots \mu_s}(x) J_{\mu_1 \dots \mu_s}(\Phi) \right]$$

- $H_{\text{vac}} = \frac{1}{2} \eta^{\mu\nu} p_\mu p_\nu$

$$\epsilon\text{-gauge inv} \quad \rightarrow \quad \partial^{\mu_1} J_{\mu_1 \dots \mu_s} = 0 \quad \text{if} \quad \square \Phi = 0$$

$$\alpha\text{-gauge inv} \quad \rightarrow \quad \eta^{\mu_1 \mu_2} J_{\mu_1 \mu_2 \dots \mu_s} - \frac{1}{2} \square J_{\mu_3 \dots \mu_s} = 0$$

- after redefinition $J_s \rightarrow$ conserved traceless Noether currents corresponding to symmetries of $\square \Phi = 0$ in R^d [Eastwood 02]
- their algebra = HS algebra of conformal spins in R^d
= HS algebra of massless spins in AdS_{d+1} [Vasiliev, Fradkin, Linetsky]

Action for h_s : [AT 02; Segal 02; Bekaert, Morad, Joung 10]

- $\log \Lambda_{UV}$ term of scalar 1-loop action $\log \det \hat{H}$

$$S[h] = \text{“Seeley coeff”} = t^0 \text{ term in } \left. \text{Tr} e^{-t\hat{H}} \right|_{t \rightarrow 0}, \quad H = H_{\text{vac}} + h$$

- inherits CHS symm: $\delta h = [H, \epsilon(x, p)]_* + \{H, \omega(x, p)\}_*$

$$\rightarrow \delta h_{\mu_1 \dots \mu_s} = \partial_{(\mu_1} \epsilon_{\mu_2 \dots \mu_s)} + \eta_{(\mu_1 \mu_2} \alpha_{\mu_3 \dots \mu_s)} + O(h)$$

- this construction can be generalized [Grigoriev, AT 16]

to curved vacuum expansion point: $H_{\text{vac}} = \frac{1}{2} g^{\mu\nu}(x) p_\mu p_\nu$

CHS as induced theory: AdS/CFT

start with free $U(N)$ scalar CFT $\int d^4x \Phi_i^* \partial^2 \Phi_i$

- tower of on-shell conserved traceless currents

$$J_s = \Phi_i^* \mathcal{J}_s \Phi_i \sim \Phi_i^* \partial_{(\mu_1} \dots \partial_{\mu_s)} \Phi_i + \dots$$

- implies infinite tower of conserved charges:

symmetries of $\square\Phi = 0 \rightarrow$ HS symmetry [Eastwood, Vasiliev]

- generating functional for correlators of currents:

add $h_s J_s$ and integrate out Φ_i

$$\Gamma[h] = N \log \det \left(-\partial^2 + \sum_s h_s \mathcal{J}_s \right), \quad \mathcal{J}_s \sim \partial^s$$

- source fields = CHS fields h_s : $\Delta(h_s) = 2 - s$

- CHS theory: gauge theory for

HS symmetries (conf Killing tensors) of $\square\Phi = 0$

(cf. Weyl gravity as gauge theory of conformal group)

$$\delta h_{\mu_1 \dots \mu_s} = \partial_{(\mu_1} \varepsilon_{\mu_2 \dots \mu_s)} + \eta_{(\mu_1 \mu_2} \alpha_{\mu_3 \dots \mu_s)} + O(h)$$

- vectorial AdS/CFT: [Klebanov, Polyakov 02]

J_s dual to massless HS fields in AdS_{d+1}

$\Gamma[h]$ should follow from Vasiliev-type theory in AdS_{d+1}

upon integrating over AdS_{d+1} fields ϕ_s with Dirichlet b.c.

$$e^{-\Gamma[h]} = \int_{\phi_s \Big|_{\partial AdS} = h_s} [d\phi_s] \exp(-NS_{HS}[\phi])$$

- full $\Gamma[h]$ is non-local and does not have CHS symmetry but its log divergent part is local and invariant:

$$\Gamma[h] \rightarrow NS_{CHS}[h] \log \Lambda_{UV} + \dots$$

$$NS_{HS}[\phi] \Big|_{\text{on-shell}} \rightarrow NS_{CHS}[h] \log \Lambda_{IR} + \dots$$

- CHS action as induced action:

$$S_{\text{CHS}} \sim \log \det \Delta(h) \Big|_{\log \Lambda_{\text{UV}}}, \quad \Delta(h) = -\partial^2 + \sum_s \mathcal{J}_s h_s$$

- familiar low-spin cases ($s = 0, 1, 2$) in covariant form

$$L = \sqrt{g} \left[g^{\mu\nu} D_\mu \Phi^* D_\nu \Phi + \left(\frac{1}{6} R + h'_0 \right) \Phi^* \Phi \right], \quad D_\mu = \partial_\mu + \frac{i}{2} h'_\mu$$

$$\begin{aligned} L &= \partial_\mu \Phi^* \partial^\mu \Phi + \sum_s h_s \Phi^* \mathcal{J}_s \Phi \\ &= \partial_\mu \Phi^* \partial^\mu \Phi + h_0 \Phi^* \Phi + i h^\mu \Phi^* \partial_\mu \Phi + \frac{1}{2} h^{\mu\nu} \partial_\mu \Phi^* \partial_\nu \Phi + \dots \end{aligned}$$

by local field redefinition ($h'_{\mu\nu} \equiv g_{\mu\nu} - \eta_{\mu\nu}$)

$$h'_0 = h_0 + \frac{1}{4} h_\mu h^\mu + \frac{1}{96} (\partial_\lambda h_{\mu\nu} \partial^\lambda h^{\mu\nu} + \dots) + \dots$$

$$h'_\mu = h_\mu + \frac{1}{2} h_{\mu\nu} h^\nu + \frac{1}{4} h_{\mu\nu} h^{\nu\lambda} h_\lambda + \dots, \quad h'_{\mu\nu} = \frac{1}{2} h_{\mu\nu} + \frac{1}{4} h_{\mu\lambda} h^\lambda_\nu + \dots$$

log divergent part of scalar log det

$$S[h'_0, h'_1, h'_2] = \int d^4x \sqrt{g} \left(h'^2_0 - \frac{1}{24} F'^2_{\mu\nu} + \frac{1}{60} C^2_{\mu\nu\lambda\rho} \right)$$

Computing CHS action as induced action

[Beccaria, Nakach, AT 16]

- 2-, 3- and 4-point vertices in CHS action

from UV pole part of scalar loop integrals with J_s insertions

- same as local limit of correlators of currents

$$\langle J_{s_1}(x_1) \dots J_{s_n}(x_n) \rangle \Big|_{x_i \rightarrow x}$$

- coupling of external CHS fields to complex scalar


$$L = -\partial_\mu \Phi^* \partial^\mu \Phi + \sum_{s=0}^{\infty} J_{\mu(s)} h^{\mu(s)}, \quad J_{\mu(s)} \equiv J_{\mu_1 \dots \mu_s}$$

$$J_{\mu(s)}(x) = \frac{i^s 2^s}{(2s)!} \sum_{k=0}^s \binom{s}{k} \binom{\frac{s+k-1}{2}}{s} G_{\mu(s)}^{(k)}(x)$$

$$G_{\mu(s)}^{(k)}(x) = (\partial - \partial')^{\mu(k)} (\partial + \partial')^{\mu(s-k)} \Phi(x) \Phi^*(x') \Big|_{x=x'}$$

$$S = \int d^4x \left(\sum_s h_s \partial^{2s} h_s + \sum_{s_i} \partial^{s_1+s_2+s_3-2} h_{s_1} h_{s_2} h_{s_3} + \sum_{s_i} \partial^{s_1+s_2+s_3+s_4-4} h_{s_1} h_{s_2} h_{s_3} h_{s_4} + \dots \right)$$

• kinetic term:




The diagram shows a circular loop with two external lines. The left external line is labeled $h(p)$ and the right external line is labeled $h(-p)$. The top arc of the loop is labeled k with an arrow pointing left, and the bottom arc is labeled $k+p$ with an arrow pointing right. To the right of the diagram is the integral expression:

$$= \int \frac{d^d k}{(2\pi)^d} \frac{N(k,p)}{k^2 (k+p)^2}$$

$\frac{1}{\varepsilon} = \frac{1}{d-4}$ UV pole part (for TT field h_s):

$$S_2 = \frac{1}{2^s (2s+1)!} \int d^4x h_{\mu(s)} \square^s h^{\mu(s)}$$

- cubic vertex: pole part of



example: 1-1-s

$$V_{\mu\nu\rho(s)} = \int \frac{d^d k}{(2\pi)^d} \frac{k_\mu (k+p_1)_\nu (k+p_1+p_2)_\rho (s)}{k^2 (k+p_1)^2 (k+p_1+p_2)^2} \Big|_{\frac{1}{\epsilon} \text{ part}}$$


$$S_3(1, 1, s) = \frac{1}{(s+2)!} \int d^4 x \left[\partial^{\rho(s)} h_\mu h^\mu h_{\rho(s)} - 2h_\mu \partial^\mu \partial_{\rho(s-1)} h_\nu h^{\nu\rho(s-1)} \right. \\ \left. - \frac{s}{2} \partial^{\rho(s-2)} \square h^\mu h^\nu h_{\mu\nu\rho(s-2)} - \frac{s}{2} \partial^{\rho(s-2)} h^\mu \square h^\nu h_{\mu\nu\rho(s-2)} \right. \\ \left. - \partial_\lambda \partial^{\rho(s-2)} h^\mu \partial^\lambda h^\nu h_{\mu\nu\rho(s-2)} \right]$$

e.g. 1-1-2 is like in Maxwell $\int d^4x \sqrt{g} g^{\mu\nu} g^{\lambda\rho} F_{\mu\lambda} F_{\nu\rho}$

$$S_3(1, 1, 2) = \frac{1}{24} \int d^4x \left[\partial_\rho h_\mu \partial_\sigma h^\mu h^{\rho\sigma} - 2 \partial_\rho h_\mu \partial^\mu h_\nu h^{\nu\rho} \right. \\ \left. + 2 h^\mu \square h^\nu h_{\mu\nu} + \partial_\lambda h^\mu \partial^\lambda h^\nu h_{\mu\nu} \right]$$

• quartic vertex:

e.g. 4-vector contact term from pole part of diagram



$\frac{1}{16} \int d^4x (h_\mu h^\mu)^2$ combining into $\int d^4x (h_0 + \frac{1}{4} h_\mu h^\mu)^2$:

contribution to 1-1-1-1 scattering cancels against h_0 exchange

• similarly for 2-2-s and 2-2-2-2 vertices, etc.

S-matrix of CHS theory in flat vacuum

[Beccaria, Nakach, AT 16]


- compute tree-level CHS 4-point amplitudes A_4 for external states = massless ($\square h_s = 0$) modes in flat space
- A_4 turns out to be zero after summation over all spin s CHS intermediate states
- this appears to be a consequence of CHS global symmetry

first illustrate this on simplest example:

scattering of external scalars via exchange of infinite tower of all CHS fields

Scalar scattering via conformal HS exchange

[Joung, Nakach, AT 15]



$$S[\Phi, h] = \int d^4x \left[\Phi^* \partial^2 \Phi + \sum_{s=0}^{\infty} h_s J_s(\Phi) \right] + S[h]$$

$$S[h] = \frac{1}{g^2} \sum_{s=0}^{\infty} \int h_s P_s \partial^{2s} h_s + \mathcal{O}(h^3)$$

- h_0 coupled to $\Phi^* \Phi$; h_μ to $i\Phi^* \partial_\mu \Phi + c.c.$; $h_{\mu\nu}$ to $T_{\mu\nu}$, etc.
- h_s exchange with propagator $\sim \frac{1}{p^{2s}}$ and p^s in the vertices:
scale invariance, no dimensional parameters

Four-scalar tree-level scattering amplitude

t-channel amplitude

$$A^{(t)}(s, t, u) = g^2 F\left(\frac{s-u}{s+u}\right), \quad F(z) \equiv \sum_{s=0}^{\infty} \left(s + \frac{1}{2}\right) P_s(z)$$

s, t, u are Mandelstam variables: $s + t + u = 0$

$P_s(z)$ – Legendre polynomial

- amplitude is scale-invariant: depends on ratios of s, t, u
- summing over spins:

$$\sum_{s=0}^{\infty} f(s) \rightarrow \sum_{s=0}^{\infty} f(s) e^{-\epsilon(s+\frac{1}{2})} \Big|_{\epsilon \rightarrow 0, \text{ fin}}$$

one finds that amplitude is δ -function in phase space

$$F(z) = \delta(z - 1)$$

Total amplitude: sum of channels

$$A_{\Phi\Phi\rightarrow\Phi\Phi} = g^2 \left[\delta\left(\frac{s}{t}\right) + \delta\left(\frac{s}{u}\right) \right]$$

in c.o.m. frame $\vec{p}_1 + \vec{p}_2 = 0 = \vec{p}_3 + \vec{p}_4$

scattering angle: $\frac{s}{t} = -(\sin^2 \frac{\theta}{2})^{-1}$, $\frac{s}{u} = -(\cos^2 \frac{\theta}{2})^{-1}$

arguments of delta-functions never vanish for real θ

$$A_{\Phi\Phi\rightarrow\Phi\Phi} = 0$$

$$A_{\Phi\Phi^*\rightarrow\Phi\Phi^*} = \frac{g^2}{2} \left[\delta\left(\frac{u}{t}\right) + \delta\left(\frac{u}{s}\right) \right] = \frac{g^2}{2} \left[\delta(\cot^2 \frac{\theta}{2}) - \delta(\cos^2 \frac{\theta}{2}) \right]$$

t-channel and s-channel contributions cancel each other

$$A_{\Phi\Phi^*\rightarrow\Phi\Phi^*} = 0$$

thus individual spin s exchange contributions are nontrivial

but total amplitude =0

- underlying HS symmetry constrains the S-matrix

(cf. integrability / hidden charges in 2d theories):

$A_4 = 0$ is implied by global part of CHS gauge symmetry:

conformal group generators plus higher spin generators

- in particular: “hyper-translations”

$$\delta\Phi = \epsilon^{m_1 \dots m_r} \partial_{m_1} \dots \partial_{m_r} \Phi$$

- fixes $A_4(\mathbf{s}, \mathbf{t}, \mathbf{u}) = k_1(\mathbf{t}, \mathbf{u}) \delta(\mathbf{s}) + k_2(\mathbf{s}, \mathbf{u}) \delta(\mathbf{t}) + k_3(\mathbf{t}, \mathbf{s}) \delta(\mathbf{u})$

- also scale invariance: $A_4(\lambda^2 \mathbf{s}, \lambda^2 \mathbf{t}, \lambda^2 \mathbf{u}) = A_4(\mathbf{s}, \mathbf{t}, \mathbf{u})$

- solution consistent with crossing and scaling symmetry

$$A_4(\mathbf{s}, \mathbf{t}, \mathbf{u}) = 0$$

★ special prescription for summation over s

with which tree-level amplitude vanishes

is thus consistent with underlying CHS symmetry

Scattering of conformal higher spin fields


[Beccaria, Nakach, AT 16]

- $s = 1$ case is standard vector but for $s \geq 2$
higher-derivative \square^s kinetic term: non-unitary theory
- definition of S-matrix: amputated Green's functions
computed with full CHS vertices and internal propagators
but with special – massless spin s – asymptotic states attached
[e.g. for $s = 2$: Weyl graviton (6 d.o.f.)
use standard helicity ± 2 gravitons as special asymptotic states]

CHS 4-particle tree level amplitude

helicities $(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$ and s, t, u ($p_i^2 = 0$ for legs)

exchange diagrams



$s = 1$ scattering: $11 \rightarrow 11$

spin s exchange: two 1-1- s vertices

and TT spin s propagator ($p_{\rho(s)} \equiv p_{\rho_1} \dots p_{\rho_s}$)

$$V_{\alpha\beta\rho(s)}(p, q) = \frac{1}{(s+2)!} \left\{ \eta_{\alpha\beta} [p_{\rho(s)} + q_{\rho(s)}] \right. \\
- \eta_{\alpha\rho_1} p_\beta p_{\rho_2} \dots p_{\rho_s} + \eta_{\beta\rho_1} q_\alpha p_{\rho_2} \dots p_{\rho_s} - \eta_{\beta\rho_1} q_\alpha q_{\rho_2} \dots q_{\rho_s} + \eta_{\alpha\rho_1} p_\beta q_{\rho_2} \dots q_{\rho_s} \\
\left. - \eta_{\alpha\rho_1} \eta_{\beta\rho_2} p_{\rho_3} \dots p_{\rho_s} p \cdot q - \eta_{\alpha\rho_1} \eta_{\beta\rho_2} q_{\rho_3} \dots q_{\rho_s} p \cdot q \right\}$$

• $s = 2$ exchange (\square^{-2}):

same as in conformal supergravity ($L = -F^2 + C^2 + \dots$)

only MHV are non-zero (++++, +++-, ... = 0)

λ	$A_s^{(2)}$	$A_t^{(2)}$	$A_u^{(2)}$
$\pm \pm \mp \mp$	0	$\frac{5}{48} \frac{s^2}{t^2}$	$\frac{5}{48} \frac{s^2}{u^2}$
$\pm \mp \mp \pm$	$\frac{5}{48} \frac{u^2}{s^2}$	$\frac{5}{48} \frac{u^2}{t^2}$	0

• $s = 4$ exchange (\square^{-4}):

again only MHV are non-zero:

λ	$A_s^{(4)}$	$A_t^{(4)}$	$A_u^{(4)}$
$\pm \pm \mp \mp$	0	$\frac{s^2 (28 s^2 + 42 s t + 15 t^2)}{80 t^4}$	$\frac{s^2 (28 s^2 + 42 s u + 15 u^2)}{80 u^4}$
$\pm \mp \mp \pm$	$\frac{u^2 (28 u^2 + 42 s u + 15 s^2)}{80 s^4}$	$\frac{u^2 (28 u^2 + 42 t u + 15 u^2)}{80 t^4}$	0

• General spin s exchange $11 \rightarrow 11$ amplitudes ($\neq 0$)

$$A_t^{(s)}(\pm \pm \mp \mp) = c_s \left(\frac{s}{t}\right)^s P_s\left(\frac{t}{s}\right), \quad A_u^{(s)}(\pm \pm \mp \mp) = c_s \left(\frac{s}{u}\right)^s P_s\left(\frac{u}{s}\right),$$

$$A_s^{(s)}(\pm \mp \mp \pm) = c_s \left(\frac{u}{s}\right)^s P_s\left(\frac{s}{u}\right), \quad A_t^{(s)}(\pm \mp \mp \pm) = c_s \left(\frac{u}{t}\right)^s P_s\left(\frac{t}{u}\right)$$

$$c_s = \frac{2s+1}{2(s-1)s(s+1)(s+2)}$$

$$P_s(x) = x^{s-2} P_{s-2}^{(4,0)}\left(\frac{x+2}{x}\right), \quad \text{order } s-2, \quad s = 2, 4, 6, \dots$$

$P_n^{(a,b)}(x)$ = Jacobi polynomials

$$P_s(x) = \sum_{j=2}^s \frac{1}{(j-2)!(j+2)!} \frac{(s+j)!}{(s-j)!} x^{s-j} \sim x^{s-2} {}_2F_1\left(2-s, s+3, 5; -\frac{1}{x}\right)$$

Sum over spins

total ++ -- amplitude: t + u-channel

$$A^{(s)} = c_s \left[\left(\frac{s}{t}\right)^s P_s\left(\frac{t}{s}\right) + \left(\frac{s}{u}\right)^s P_s\left(\frac{u}{s}\right) \right]$$

$$A^{(s)} = \sigma_s(x) + \sigma_s(-1-x), \quad \sigma_s(x) = c_s x^{-s} P_s(x), \quad x = \frac{t}{s}$$

use generating function for Jacobi polynomials $P_{s-2}^{(4,0)}$

$$\sum_{s=2}^{\infty} x^{-s} P_s(x) z^{s-2} = \frac{1}{x^2} \frac{16}{\sqrt{z^2 - \frac{2z(x+2)}{x} + 1} \left(\sqrt{z^2 - \frac{2z(z+2)}{z} + 1} - z + 1 \right)^4}$$

$$\begin{aligned} \sigma(x) &= \sum_{s=2,4,6,\dots}^{\infty} \sigma_s(x) = \lim_{z \rightarrow 1} \sum_{s=2,4,6,\dots}^{\infty} c_s x^{-s} P_s(x) z^{s-2} \\ &= \frac{1}{8} \left[-2x + 2(x+1)x \log\left(\frac{1}{x} + 1\right) - 1 \right]. \end{aligned}$$

total amplitude is zero as in scalar scattering case

$$A(x) = \sum_{s=2,4,6,\dots}^{\infty} A^{(s)}(x) = \sigma(x) + \sigma(-1-x) = 0$$

$s = 2$ scattering

- $+2+2 \rightarrow +2+2$: contribution from from $s > 2$ exchanges:
t-channel $++ \rightarrow ++$ or $+- -$ MHV

$$A_{++;++}(t, \theta) = \frac{s^4}{t^4} \sum_{s=4,6,\dots} (s + \frac{1}{2}) F^{(s)} t^2 P_{s-4}^{(8,0)}(\cos \theta)$$

explicit computation gives for full (t + u- channel) amplitude

$$A^{(s)} = c_s s^2 \left[\left(\frac{s}{t}\right)^{s-2} P_s\left(\frac{t}{s}\right) + \left(\frac{s}{u}\right)^{s-2} P_s\left(\frac{u}{s}\right) \right]$$
$$P_s(x) = x^{s-2} P_{s-4}^{(8,0)}\left(\frac{x+2}{x}\right), \quad c_s = \frac{9}{32} \frac{2s+1}{(s-3)\dots(s+4)}$$

- sum over spins:

$$\sigma(x) = \sum_{s=4,6,\dots}^{\infty} \sigma_s(x) = \lim_{z \rightarrow 1} \sum_{s=4,6,\dots}^{\infty} c_s x^{-(s-2)} P_s(x) z^{s-4}$$
$$= \frac{1}{4320} \left[60 (x+1)^3 x^3 \log\left(\frac{1}{x} + 1\right) - 60 x^5 - 150 x^4 - 110 x^3 - 15 x^2 + 3 x - 1 \right]$$

- total $s > 2$ exchange vanishes: t- and u- channels cancel

$$\sigma(x) + \sigma(-1 - x) = 0$$

- contribution of $s = 0, 2$ exchanges + 2222 contact vertex

$$A_{++++}^{0,s} = \frac{s^2}{18432}, \quad A_{++++}^{0,t} = \frac{t^2 u^4}{2048 s^4}, \quad A_{++++}^{0,u} = \frac{t^4 u^2}{2048 s^4},$$

$$A_{++++}^{2,s} = \frac{s^2 + 6 s t + 6 t^2}{92160}, \quad A_{++++}^{2,t} = \frac{u^2 (2 s^4 - 10 s^3 t + 33 s^2 t^2 - 24 s t^3 + 3 t^4)}{30720 s^4}$$

$$A_{++++}^{2,u} = \frac{t^2 (2 s^4 - 10 s^3 u + 33 s^2 u^2 - 24 s u^3 + 3 u^4)}{30720 s^4}$$

$$A_{++++}^{\text{contact}} = -\frac{s^6 - s^5 t + 26 s^4 t^2 + 63 s^3 t^3 + 54 s^2 t^4 + 27 s t^5 + 9 t^6}{7680 s^4}$$

non-trivial cancellation: total 2222 amplitude = 0

$$A^{0,s} + A^{0,t} + A^{0,u} + A^{2,s} + A^{2,t} + A^{2,u} + A^{\text{contact}} = 0$$

- similar cancellation checked for 1122 amplitude
 - conjecture: full massless-state CHS S-matrix is trivial
 - this should follow from underlying global CHS symmetry
- HS charges \rightarrow triviality of S-matrix (cf. Coleman-Mandula)

CHS symmetries

$$h(x, p) \equiv h_{\mu_1 \dots \mu_s}(x) p^{\mu_1} \dots p^{\mu_s}$$

$$f(x, p) \star g(x, p) = f(x, p) e^{\frac{i}{2}(\overleftarrow{\partial}_x \cdot \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \cdot \overrightarrow{\partial}_x)} g(x, p)$$

- diff and algebraic symm of scalar-CHS system [Segal 02]

$$\delta_\epsilon h(x, p) = (p \cdot \partial_x) \epsilon(x, p) - \frac{i}{2} [h(x, p), \epsilon(x, p)]_\star$$

$$\delta_\alpha h(x, p) = (p^2 - \frac{1}{4} \partial_x^2) \alpha(x, p) - \frac{1}{2} \{h(x, p), \alpha(x, p)\}_\star$$

$$\delta_{\epsilon+i\alpha} \Phi(x) = e^{-\frac{i}{2} \partial_{x'} \cdot \partial_p} [\epsilon(x, p) + i\alpha(x, p)] \Phi(x') \Big|_{x=x', p=0}$$

$$\delta h = \delta_0 h + \delta_1 h, \quad \delta_0 h_s \sim \partial \epsilon_{s-1} + \eta \alpha_{s-2}$$

- global symmetry from: $\delta_1 h \sim \epsilon \partial h + \partial \epsilon h + \dots$ for special ϵ

- spin s field transforms in terms of $s' < s$ fields

$$\delta_1 h_0 \sim \sum_k \epsilon^{\mu(k)} \partial_{\mu(k)} h_0, \quad \delta_1 h^\rho \sim \sum_k [\epsilon^{\rho\mu(k)} \partial_{\mu(k)} h_0 + \epsilon^{\mu(k)} \partial_{\mu(k)} h^\rho]$$

$$\delta_1 h^{\rho\sigma} \sim \sum_k [\epsilon^{\rho\sigma\mu(k)} \partial_{\mu(k)} h_0 + \epsilon^{\mu(k)(\rho} \partial_{\mu(k)} h^{\sigma)} + \frac{1}{2!k!} \epsilon^{\mu(k)} \partial_{\mu(k)} h^{\rho\sigma}]$$

- constraints on amplitudes as in external scalar scattering case

CHS fields in curved background

Expansion near vacuum with non-trivial $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$:

- Weyl-invariant quadratic action known for $s = 1$ and $s = 2$
- $s > 2$: kinetic operator $\mathcal{O}_s = \nabla^{2s} + \dots$ – diff and Weyl inv

but to be consistent with CHS gauge symm.:

$g_{\mu\nu}$ should solve Bach eqs $(\nabla^\mu \nabla^\nu + \frac{1}{2} R^{\mu\nu}) C_{\lambda\mu\nu\rho} = 0$

- \mathcal{O}_s simplifies/factorizes on conf-flat background:

explicitly known on S^4 or AdS_4 [AT 13; Metsaev 14; Nutma, Taronna 14]

and $S^1 \times S^3$ [Bekaert, Beccaria, AT 14]

- quantum consistency? anomalies?

conformal \rightarrow Weyl symmetry: $g'_{mn} = \lambda^2(x) g_{mn}$

Weyl anomaly: $T_m^m = -a R^* R^* + c C^2$

Weyl gravity is anomalous: $a_2 = \frac{87}{20}$, $c_2 = \frac{199}{30}$

- one way to cancel anomaly – add fermions: **supersymmetry**

$N = 4$ conformal supergravity + 4 $N = 4$ Maxwell multiplets

is anomaly free: $a = c = 0$ [Fradkin, AT 82]

- alternative: sum over infinite number of CHS contributions

- CHS fields with $s > 2$:

a_s : enough to know partition function on S^4

c_s : need to know \mathcal{O}_s on Ricci-flat background

CHS partition function on S^4

- Maxwell theory on S^4 ($R = 12$, $r = 1$)

$$Z_1 = \left[\frac{\det \Delta_0(0)}{\det \Delta_{1\perp}(3)} \right]^{1/2}, \quad \Delta_s(M^2) \equiv -\nabla_s^2 + M^2$$

- Weyl graviton: $C^2 \rightarrow \frac{1}{2} h \Delta_{2\perp}(2) \Delta_{2\perp}(4) h$

$$Z_2 = \left[\frac{\det \Delta_{1\perp}(-3)}{\det \Delta_{2\perp}(2)} \right]^{1/2} \left[\frac{\det \Delta_0(-4)}{\det \Delta_{2\perp}(4)} \right]^{1/2}$$

- CHS operator: factorization into “partially-massless” ops

$$\mathcal{O}_s = \nabla^{2s} + \dots = \prod_{k=0}^{s-1} \Delta_{s\perp}(M_{s,k}^2), \quad M_{s,k}^2 = 2 + s - k - k^2$$

- get simple generalization of flat-space Z

$$Z(S^4) = \prod_{s=1}^{\infty} Z_s, \quad Z_s = \prod_{k=0}^{s-1} Z_{s,k}, \quad Z_{s,k} = \left[\frac{\det \Delta_{k\perp}(M_{k,s}^2)}{\det \Delta_{s\perp}(M_{s,k}^2)} \right]^{1/2}$$

$$\ln Z = -B_4 \ln \Lambda_{\text{UV}} + \dots, \quad B_4 = \int d^4x \sqrt{g} b_4 \Big|_{S^4} = -a_s$$

- summing contributions of 2nd order operators [AT 13]

$$\begin{aligned} a_s &= \sum_{k=0}^{s-1} \left(a[\Delta_{s\perp}(2 + s - k - k^2)] - a[\Delta_{k\perp}(2 + k - s - s^2)] \right) \\ &= \frac{1}{180} \nu^2 (14\nu + 3), \quad \nu = s(s + 1) \end{aligned}$$

- same coefficient found via massless HS AdS_5 relation

[Giombi, Klebanov, Pufu, Safdi, Tarnapolsky 13]

$$\ln \frac{Z_s^{(-)}}{Z_s^{(+)}} = \ln Z_s = a_s \ln \Lambda_{\text{IR}} + \dots, \quad \text{vol}(AdS_5) \sim \ln \Lambda_{\text{IR}}$$

- with $e^{-\epsilon(s+\frac{1}{2})}$ regularization prescription for \sum_s consistent with CHS symmetries get

$$\sum_{s=1}^{\infty} a_s = 0$$

- finite parts cancel too: $Z(S^4) = 1$

[Giombi, Klebanov, Safdi 14; Beccaria, AT 15]

Ricci-flat background

- Maxwell vector: $(\Delta_1)_{mn} = -(\nabla^2)_{mn} + R_{mn}$, $\Delta_0 = -\nabla^2$

$$Z_1 = \left[\frac{(\det \Delta_0)^2}{\det \Delta_1} \right]^{1/2}$$

- Weyl graviton: 4-th order operator factorizes: [\[Fradkin, AT 82\]](#)

square of Einstein op. $(\Delta_2)_{mn,kl} = -(\nabla^2)_{mn,kl} - 2C_{mknl}$

$$Z_2 = \left[\frac{(\det \Delta_1)^3}{(\det \Delta_2)^2} \right]^{1/2}$$

- assume factorization of \mathcal{O}_s true also for $s > 2$:
 s factors of “massless” spin s 2nd-order operator

$$Z_s = \left[\frac{(\det \Delta_{s-1})^{s+1}}{(\det \Delta_s)^s} \right]^{1/2}, \quad \Delta_s = -\nabla^2 - s(s-1)C \dots$$

same structure as in flat space but with covariant operators Δ_s

- from Seeley coefficients for Δ_s get [AT 13]

$$c_s - a_s = \frac{1}{720} \nu_s (15\nu_s^2 - 45\nu_s + 4), \quad \nu_s = s(s+1)$$

with same summation over spins prescription

$$\sum_{s=1}^{\infty} (c_s - a_s) = 0$$

- then a- and c- anomalies or UV ∞ appear to vanish:
suggests novel mechanism of UV finiteness due to
summation of ∞ number of bosonic fields (cf. string theory)

- But $\sum_s c_s = 0$ remains a conjecture:

- $\mathcal{O}_{s>2}$ does not factorize on $R_{mn} = 0$ backgr [Nutma, Taronna 14]

but obstruction to factorization $\sim \nabla.C_{\dots}$ should not change c_s

- CHS kin. op. does not diagonalize on $R_{mn} = 0$ backgr:
mixing terms [Grigoriev, AT 16] contribute to c_s [Beccaria, AT 17]

Curved space background: spin 1 – 3 mixing

[Beccaria, AT 17]

• flat space:

$$S_0 = \int d^4x \Phi^* \partial^2 \Phi, \quad \partial^{a_1} J_{a_1 \dots a_s} = 0, \quad J_{a_1 \dots a_s}^{a_1} = 0$$

$$J_a = i \Phi^* \partial_a \Phi + c.c.$$

$$J_{ab} = \Phi^* \partial_a \partial_b \Phi - 2 \partial_a \Phi^* \partial_b \Phi + \frac{1}{2} \eta_{ab} \partial^c \Phi^* \partial_c \Phi + c.c.$$

$$J_{abc} = i \left[\Phi^* \partial_a \partial_b \partial_c \Phi - 9 \partial_{(a} \Phi^* \partial_b \partial_{c)} \Phi + 3 \eta_{(ab} \partial^p \Phi^* \partial_p \partial_{c)} \Phi \right] + c.c.$$

adding interaction with background fields:

$$S_{int} = \sum_s \int d^4x h^{a_1 \dots a_s}(x) J_{a_1 \dots a_s}$$

inv under $\delta h_{a_1 \dots a_s} = \partial_{(a_1} \epsilon_{a_2 \dots a_s)} + \eta_{(a_1 a_2} \alpha_{a_3 \dots a_s)}$ mod $\partial^2 \Phi$ terms

extended off shell if transform Φ and add terms linear in h_s

• curved space:

$$S_0 = \int d^4x \sqrt{g} \Phi^* \left(-\nabla^2 + \frac{1}{6} R \right) \Phi$$

$$S_{int} = \sum_s \int d^4x \sqrt{g} h^{a_1 \dots a_s}(x) J_{a_1 \dots a_s}$$

- require $\nabla^{a_1} J_{a_1 \dots a_s} = 0$, $J^{a_1}_{a_1 \dots a_s} = 0$

then will have inv under backgr-cov gauge transfs

$$\delta h_{a_1 \dots a_s} = \nabla_{(a_1} \epsilon_{a_2 \dots a_s)} + g_{(a_1 a_2} \alpha_{a_3 \dots a_s)}$$

- require also Weyl inv w.r.t. backgr metric: $\omega = \omega(x)$

$$\delta_{\text{W}} g_{ab} = 2\omega g_{ab}, \quad \delta_{\text{W}} \Phi = -\omega \Phi, \quad \delta_{\text{W}} h_{a_1 \dots a_s} = 2(s-1)\omega h_{a_1 \dots a_s}$$

- such covariant currents exist for $s = 1$ and $s = 2$:

$$J_a = i (\Phi^* \nabla_a \Phi - \nabla_a \Phi^* \Phi), \quad \nabla^a J_a = 0$$

$$J_{ab} = \frac{6}{\sqrt{g}} \frac{\delta S_0}{\delta g^{ab}} = (\Phi^* \nabla_a \nabla_b \Phi - 2 \nabla_a \Phi^* \nabla_b \Phi + c.c.) \\ + g_{ab} \nabla_c \Phi^* \nabla^c \Phi - (R_{ab} - \frac{1}{6} g_{ab} R) \Phi^* \Phi$$

- but $s \geq 3$ cases are different:

$\nabla^{a_1} J_{a_1 \dots a_s} \neq 0$ – given by terms with lower-rank J_s

- $s = 3$: **unique** traceless current with Weyl-inv $S_{int} = \int h_3 J_3$:

$$J_{abc} = i \left[\Phi^* \nabla_{(a} \nabla_b \nabla_{c)} \Phi - 9 \nabla_{(a} \Phi^* \nabla_b \nabla_{c)} \Phi + 3 g_{(ab} \nabla^p \Phi^* \nabla_p \nabla_{c)} \Phi \right. \\ \left. + 2 g_{(ab} \Phi^* \nabla^2 \nabla_{c)} \Phi + \frac{1}{2} g_{(ab} R \Phi^* \nabla_{c)} \Phi - 7 R_{(ab} \Phi^* \nabla_{c)} \Phi \right] + c.c.$$

- J_3 conserved only in conformally-flat background:

$$\nabla_a J^{abc} = 8 C^{pbcq} \nabla_{(p} J_{q)} + 32 \nabla_{(p} C^{pbcq} J_{q)}$$

- 1+3 action $S_{int} = \int d^4x \sqrt{g} (h^a J_a + h^{abc} J_{abc})$

is invariant under $\delta h_a = \partial_a \epsilon$ and combined transformations

$$\delta h_{abc} = \nabla_{(a} \epsilon_{bc)}, \quad \delta h_a = -8 C_{abcd} \nabla^d \epsilon^{bc} + 24 \nabla^d C_{abcd} \epsilon^{bc}$$

- to make invariance manifest (off-shell):

need also to transform Φ and add $h_1 h_3 + \dots$ terms in S_{int}

(manifest spin 1 invariance: $\nabla_a \Phi \rightarrow D_a \Phi = \nabla_a \Phi + i h_a \Phi$)

- induced action inv under h -gauge transf

$$e^{-\Gamma(h)} = \int d\Phi e^{-S(\Phi, h; g)}, \quad \Gamma(h) = S(h) \log \Lambda_{UV} + \dots$$

- non-linear $h_s h_{s'} + \dots$ terms produce contact terms in generating functional for correlators of currents: absent in correlators $\langle J(x_1) \dots J(x_n) \rangle$ at separated points but contributing to local UV singular part, i.e. to induced action

- need contact terms to get e.g. covariant spin 1 + 2 action

$$S = \int d^4x \sqrt{g} \left(-\frac{1}{12} F_{ab}^2 + \frac{1}{120} C_{abcd}^2 \right)$$

- expansion near g_{ab} :

$$\int d^4x \sqrt{g} C_{abcd}^2 \rightarrow \int d^4x \sqrt{g} \left[B_{ab}(g) h^{ab} + h^{ab} \mathcal{O}_{abcd}(g) h^{cd} + \dots \right]$$

$\mathcal{O}_4 = \nabla^4 + \dots$ is gauge-inv $\delta h_{ab} = \nabla_{(a} \epsilon_{b)}$ if

$$B_{ab} = \left(\nabla^p \nabla^q + \frac{1}{2} R^{pq} \right) C_{apqb} = 0$$

- expansion of S in h_s : manifest reparam and Weyl inv

$$S(g, h) = S^{(0)}(g) + S^{(1)}(g, h) + S^{(2)}(g, h) + \dots$$

$$S^{(1)} = \int B_{(s)}(g) h^{(s)}, \quad S^{(2)} = \int h^{(s)} \mathcal{O}_{s,s'}(g) h^{(s')}$$

- gauge invariance if $\langle J_{(s)} \rangle_{UV} \sim B_{(s)}(g) = 0$

Weyl-inv + $\nabla^a B_{a\dots} = 0 \rightarrow$ true for Bach-flat g [Grigoriev, AT]

Quadratic part of spin 1 + 3 induced action

$$S^{(2)} = S_{11} + S_{13} + S_{33}$$

$$L_{11} = h^a \langle J_a J_b \rangle_{UV} h^b = -\frac{1}{6} F_{ab}^2$$

$$L_{33} = h_3 \mathcal{O}_6 h_3: \quad \mathcal{O}_6 \text{ from } \langle J_{abc} J_{pqr} \rangle_{UV} + \text{contact term}$$

$$L_{13} = h^a \langle J_a J_{bcd} \rangle_{UV} h^{bcd} + \text{contact term}$$

final result for the mixing term:

$$L_{13} = 8 F^{ab} \left[C_a^{cdp} \nabla_p h_{bcd} + \left(\nabla_a R^{cd} - \nabla^c R_a^d \right) h_{bcd} \right]$$

Weyl-inv; vanishes for conformally-flat Einstein space g_{ab}

- in Bach-flat case: e.g. Einstein background $R_{ab} = \frac{1}{4} R g_{ab}$


$$S^{(2)} = \int d^4x \sqrt{g} \left[-\frac{1}{12} F_{ab}^2 + 8 C^{abcd} F_{ap} \nabla_d h^p_{bc} + h_3 \mathcal{O}_6 h_3 \right]$$

- invariant under spin 3 gauge transformations

$$\delta h_{abc} = \nabla_{(a} \epsilon_{bc)} , \quad \delta h_a = -8 C_{abcp} \nabla^p \epsilon^{bc}$$

- ϵh_3 term in variation of S_{13} is order CC :

$h_3 \mathcal{O}_6 h_3$ inv by itself only to 1st order in C [Nutma, Taronna 14]



linear in curvature terms in \mathcal{O}_6 can be found from UV part of $h_2 h_3 h_3$ 1-loop scalar diagrams: $\langle J_{abc} J_{pqr} J_{mnn} \rangle_{\text{UV}} + \text{contact terms}$


Spin 1–3 mixing term contribution to UV divergences

$$\Gamma = -\log Z = -\log \Lambda_{\text{UV}} \int d^4x \sqrt{g} b_4(x) + \text{finite}$$

$$b_4 = -a R^* R^* + c C^2$$

• conf flat background: no mixing terms, \mathcal{O}_s factorize and get

$$a_s = \frac{1}{720} \nu_s (3 \nu_s + 14 \nu_s^2), \quad \nu_s \equiv s(s+1)$$



- ignoring mixings and assuming that factorization holds also in Ricci-flat case [AT 13]

$$c_s \equiv c_{ss} = \frac{1}{720} \nu_s (29 \nu_s^2 - 42 \nu_s + 4)$$

- need to add mixing terms contribution to C^2 div:

example of 1–3 sector: $c_1 = \frac{1}{10}$, $c_3 = \frac{919}{15}$

$$L = h_1(\nabla^2 + \dots)h_1 + C\nabla h_1\nabla h_3 + h_3(\nabla^6 + \dots)h_3$$

1-loop diagram gives **non-trivial** contribution:

$$L_{UV} = c_{13}C_{abcd}C^{abcd} \log \Lambda_{UV} , \quad c_{13} = \frac{392}{5}$$

- need to find all mixing terms to decide if $\sum_{s,s'} c_{ss'} = 0$

Conclusions

- theories with infinite number of massless higher spin fields:
importance of definition of quantum theory
consistent with underlying symmetries
- remarkable simplifications due to large HS symmetry:
- 1-loop $Z = 1$ on R^4 ($\sum_s \nu_s = 0$) and S^4 ($\sum_s a_s = 0$)
- vanishing of scattering amplitudes with CHS exchange:
triviality of S-matrix implied by conformal HS symmetry
- intricate structure of interacting induced CHS action
mixing terms in non-trivial background to be understood \rightarrow
cancellation of c-anomalies $\sum_s c_s = 0$ remains to be proved